

*Pogue et al.*  
*Serial No.: 10/061,216*  
*Page 2 of 5*

**AMENDMENTS TO THE CLAIMS:**

This listing of claims will replace all prior versions and listing of the claims in the application:

**LISTING OF THE CLAIMS:**

Claims 1-69 (Previously canceled)

Claim 70 (previously presented) A method of isolating a virus, comprising:

- (a) Homogenizing virus-containing plant tissue in  $\text{Na}_2\text{S}_2\text{O}_5$ ;
- (b) Straining the homogenate to obtain green juice;
- (c) Adjusting the pH of the green juice to 5.0 with acid;
- (d) Heating the green juice to about 47° C for a period of about 5 minutes followed by cooling to about 5° C;
- (e) Centrifuging the green juice at about 6000 x g for about 3 minutes to obtain a supernatant and pellet;
- (f) Precipitating the supernatant in polyethylene glycol and NaCl to obtain a precipitate;
- (g) Resuspending the precipitate in water at a concentration of about 1 mg per ml;
- (h) Extracting the precipitate in chloroform and butanol and centrifuging the extract;
- (i) Recovering and lyophilizing the aqueous phase of the centrifuged material;
- (j) Resuspending the lyophilized material at a concentration of about 5 to about 10 mg per ml water.

Claims 71-78 (Canceled)

*Pogue et al.*  
*Serial No.: 10/061,216*  
*Page 3 of 5*

**Claim 79 (New):** A method for extracting a virus from plant tissue, comprising the steps of:

homogenizing virus-containing plant tissue to obtain green juice;  
adjusting the pH of the green juice to 5.0 with acid;  
heating the green juice to about 47° C;  
cooling the green juice;  
centrifuging the green juice at about 6000 x g to obtain a supernatant and pellet;  
precipitating the supernatant in polyethylene glycol and NaCl to obtain a precipitate;  
re-suspending the precipitate in water at a concentration of about 1 mg per ml;  
extracting the precipitate in chloroform and butanol and centrifuging the extract;  
recovering and lyophilizing the aqueous phase of the centrifuged material;  
re-suspending the lyophilized material at a concentration of about 5 to about 10 mg per ml water.

**Claim 80 (New)** A method as set forth in claim 79, wherein said homogenizing step includes homogenizing the virus-containing plant tissue in  $\text{Na}_2\text{S}_2\text{O}_5$ .

**Claim 81 (New)** A method as set forth in claim 80, further comprising the step after said homogenizing step of straining the homogenate to obtain the green juice.

**Claim 82 (New)** A method as set forth in claim 81, wherein in said heating step the green juice is heated for a period of about 5 minutes.

*Pogue et al.*  
*Serial No.: 10/061,216*  
*Page 4 of 5*

Claim 83 (New) A method as set forth in claim 82, wherein in said cooling step, the green juice is cooled to about 5° C;

Claim 84 (New) A method as set forth in claim 83, wherein in said centrifuging step, the green juice is centrifuged for about 3 minutes.